

Growforce™ Cal Bor

Readily available nitrate free calcium and boron solution

Guaranteed Analysis

Calcium (Ca)5.0%
Boron (B)1.7%

Derived from: Calcium acetate and boric acid

Shake well before using.

KEEP OUT OF REACH OF CHILDREN



DANGER

May damage fertility or the unborn child. Causes eye irritation.

PRECAUTIONARY STATEMENTS

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves. Wash hands thoroughly after handling.

FIRST AID

IF EXPOSED OR CONCERNED: Get medical advice/attention.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

STORAGE AND DISPOSAL

Store locked up. Dispose of contents/container in accordance with Federal, state and local laws and regulations.

COMPATIBILITY

Growforce Cal Bor is compatible with a wide range of commonly used agricultural chemicals. When mixing with other chemicals, always mix a small quantity and check for precipitate (fall out) in the mixture. If precipitation occurs, DO NOT proceed with application. If the mixture does not precipitate and it's the first time you have used this combination of products, test a small area of the crop prior to treatment of the whole crop.

WARNING: This fertilizer carries added Boron and is intended for use only on directed crops. Its use on other crops or under conditions other than those recommended may result in serious crop injury.

DIRECTIONS FOR USE

Foliar and/or Fertigation Application:

Crop	Rate/ Acre	Comments
Apples	1 – 1.5 qts (Foliar)	Apply to newly hardened spring flush or during active growing period and post-harvest.
Avocados	1 – 1.5 qts (Foliar) 4 – 6 qts (Fertigation)	Apply to recently hardened spring flush or during active growing period and post-harvest
Canola, Corn	1 – 2 qts (Foliar)	Best applied at the late cabbage stage, can also be used at other stages
Grain legumes, Triticale, Lupins and Wheat, Cereal and Forage Legumes	1 – 2 qts (Foliar)	Apply at 3 – 4 leaf stage. For forages, apply as a stubble spray after cutting and when growth is observed
Cotton	1 – 2 qts (Foliar)	Apply from first flowering until 2 weeks before harvest.
Citrus	1 – 2 qts (Foliar)	Apply to newly hardened spring flush or during active growing period and post-harvest.
Cucurbits	1 – 2 qts (Foliar)	Apply at regular intervals post flowering until harvest.
Nut Crops; Almonds	1 – 2 qts (Foliar)	Apply at early stages of growing season when sufficient leaf cover present. Apply with compatible crop protection sprays.
Potatoes	1.5 – 3 qts (Foliar)	Regular applications from hooking until spray-off.
Stone Fruit	3 – 4 qts (Fertigation)	DO NOT apply on foliage. Best applied through soil.
Strawberries	1 – 2 qts (Foliar)	Spray from early vegetative until final harvest. DO NOT spray if fruit to be harvested within one week.
Vegetables	1 – 2 qts (Foliar)	Apply from early vegetative until a week before harvest.
Vines	1 – 2 qts (Foliar)	Apply when shoots 10 cm long. Second application: 5% flowering. Bunch finish for table grapes

WARRANTY DISCLAIMER

Vivid Life Sciences, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. Vivid Life Sciences, LLC makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

REV032619

NET CONTENTS: 2.5 GAL (9.46 L)
WEIGHT PER GALLON: 10.18 lbs. (4.62 Kg)
NET WEIGHT: 25.45 lbs. (11.54 Kg)

Manufactured for and Guaranteed By:

Vivid Life Sciences, LLC, P.O. Box 41632, Minneapolis, MN 55441